

Monopoles, confinement and the photon propagator in QED₃

M. N. Chernodub^{a *}, E.-M. Ilgenfritz^{b †} and A. Schiller^{c ‡}

^aITEP, B.Chermushkinskaya 25, Moscow, 117259, Russia and
Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan

^bResearch Center for Nuclear Physics, Osaka University, Osaka 567-0047, Japan

^cInstitut für Theoretische Physik, Universität Leipzig, D-04109 Leipzig, Germany

We study the lattice gauge boson propagator of 3D compact QED in Landau gauge at zero and non-zero temperature. Non-perturbative effects are reflected by the generation of a mass m , by an anomalous dimension α and by the photon wave function renormalisation Z . These effects can be attributed to monopoles: they are absent in the propagator of the regular part of the gauge field. The rôle of Gribov copies is carefully investigated.

Three-dimensional compact electrodynamics (cQED₃) shares two essential features with QCD, confinement [1] and chiral symmetry breaking [2]. Confinement of electrically charged particles is caused by a plasma of monopoles which emerge due to the compactness of the gauge field. Recently we have interpreted the deconfinement phase transition in cQED₂₊₁ from the monopole point of view and have shown that the transition is independent of the strength of external fields [3]. In Ref. [4] we have demonstrated how the confinement property is manifest in the gauge boson propagator of this theory and how the propagator changes at the deconfinement temperature. We have found that an anomalous dimension α appears which modifies the momentum dependence, apart from the generation of a mass m , which can be well described by Polyakov's theory [1]. We have shown that all nontrivial effects originate from the singular fields of the monopoles and disappear with the formation of dipoles.

Here we report on an extension of this study to the case of $T = 0$ and a careful investigation on the severity of the Gribov copy problem. At $T \neq 0$, it is important to realize that more structure functions are necessary for a full description. We found that one of the finite T propagators

is extremely gauge fixing sensitive. For our lattice study we have adopted the Wilson action, $S[\theta] = \beta \sum_p (1 - \cos \theta_p)$, where θ_p is the $U(1)$ field strength tensor corresponding to the compact link field θ_l . The lattice coupling β is related to the lattice spacing a and the continuum coupling constant g_3 of the 3D theory, $\beta = 1/(a g_3^2)$. The Landau gauge is defined as the maximum of a functional $\mathcal{F} = \sum_l \cos \theta_l^G$ with respect to gauge transformations G . For the Monte Carlo algorithm (combining local and global updates) and the gauge fixing procedure see Ref. [5]. Simulations for $T = 0$ have been performed on a 32^3 lattice, those for $T > 0$ on a $32^2 \times 8$ lattice.

We have studied the gauge boson propagator in Landau gauge in lattice momentum space $\vec{k}$ where it is defined as $D_{\mu\nu}(\vec{p}) = \langle \tilde{A}_{\vec{k},\mu} \tilde{A}_{-\vec{k},\nu} \rangle$ in terms of the Fourier transformed gauge potential $\tilde{A}_{\vec{k},\mu}$ for $p_\nu = (2/a) \sin(2\pi k_\nu/L_\nu)$. The gauge field in lattice position space is taken as $A_{\vec{n}+\frac{1}{2}\vec{\mu},\mu} = \sin \theta_{\vec{n},\mu}/(g_3 a)$. The most general tensor structure of $D_{\mu\nu}$ at $T = 0$ is given by

$$D_{\mu\nu}(\vec{p}) = P_{\mu\nu}(\vec{p})D(p^2) + (p_\mu p_\nu)/p^2 F(p^2)/p^2 \quad (1)$$

with the 3-dimensional transverse projection operator $P_{\mu\nu}(\vec{p}) = \delta_{\mu\nu} - (p_\mu p_\nu)/p^2$. If the Landau gauge is fulfilled exactly, $F(p^2) \equiv p_\mu D_{\mu\nu} p_\nu = 0$. The transverse and longitudinal propagators, D and F , are extracted by projection and are found approximately rotationally invariant or vanishing.

For $T > 0$ the propagator lacks $O(3)$ rotational

*M. N. Ch. is supported by JSPS Fellowship P01023.

†E.-M. I. thanks for the support by the Ministry of Education, Culture and Science of Japan (Monbu-Kagaku-sho) and the hospitality experienced at RCNP.

‡Presented by A. S. at Lattice'02.

symmetry. Thus, we have to consider two scalar functions $D_{T/L}$ using the 2-dimensional transverse/longitudinal projection operators $P_{\mu\nu}^{T/L}$:

$$D_{\mu\nu}(\vec{p}) = P_{\mu\nu}^T D_T + P_{\mu\nu}^L D_L + (p_\mu p_\nu)/p^2 F/p^2. \quad (2)$$

These projectors are given as follows ($i, j = 1, 2$; $\mathbf{p}^2 = p_i p_i$): $P_{ij}^T = \delta_{ij} - (p_i p_j)/\mathbf{p}^2$, $P_{33}^T = P_{3i}^T = P_{i3}^T = 0$ and $P_{\mu\nu}^L = P_{\mu\nu} - P_{\mu\nu}^T$. The functions $D_T(|\mathbf{p}|, p_3)$ and $D_L(|\mathbf{p}|, p_3)$ can be extracted from $D_{\mu\nu}$. In the static limit, $p_3 = 0$, we have $D_L(|\mathbf{p}|, 0) \equiv D_{33}(|\mathbf{p}|, 0)$, *i.e.* D_L contains only temporal gauge degrees of freedom directly exhibiting confinement, while D_T contains the spatial gauge degrees of freedom.

In order to discuss the form of the propagator from the monopole plasma point of view, we decompose the gauge fields links into singular (monopole) and regular (photon) contributions, $\theta_{\vec{n}, \mu} = \theta_{\vec{n}, \mu}^{phot} + \theta_{\vec{n}, \mu}^{mono}$ [6] (see also [5]). Once the decomposition of the link angles is done, one is free to define the corresponding propagators in position space, transform to momentum space, and to define the decomposition of D or $D_{L/T}$ (and F) into monopole, photon and mixed contributions.




Figure 1. The propagator $p^2 D$ and its contributions, as well as the (vanishing) propagator F vs. p^2 at $\beta = 1.8$. Only part of the data points averaged over the same p^2 are plotted.

In Fig. 1 we show for $T = 0$ at $\beta = 1.8$ the measured transverse propagator D and its different components in lattice momentum space. To create the best Landau gauge realization, we have evaluated $N_G = 20$ Gribov copies, independent maxima of $\mathcal{F}$ obtained from maximizing random

gauge transforms of the original configuration. The monopole part of the transversal propagator D^{mono} reaches a maximum in the low momentum region before it drops towards $p^2 = 0$. The regular (photon) part is singular at $p^2 \rightarrow 0$ like $D^{phot} \sim 1/p^2$, while the full transversal propagator is not. $p^2 D^{phot}$ is flat independently of β . As expected, the longitudinal part F vanishes within errors.

Following [4] we describe D by the function

$$D(p^2) = (Z/\beta) m^{2\alpha} / [p^{2(1+\alpha)} + m^{2(1+\alpha)}] + C. \quad (3)$$

The photon part D^{phot} is fitted using (3) with $\alpha = m = 0$. At finite T , the propagator data for D_L and D_T are analyzed for $p_3 = 0$, as a function of $\mathbf{p}^2$ using the same fit function (3). We have investigated the Gribov copy dependence of D and the finite T propagators D_L and D_T . In this respect, the zero temperature D behaves similarly as D_L . Some results are summarized in Fig. 2 as




Figure 2. Fitted α and m vs. N_G for D_T and D_L at different β 's and updates. For D_L at $\beta = 2.6$ we have $\alpha = 0$ and $m = 0$.

β values below and above the phase transition. After a few Gribov copy attempts N_G all fit parameters of D_L are almost insensitive to N_G . The results at large N_G are independent of whether the total Monte Carlo algorithm includes global updates or not [5] (see the label “local ” for the latter case).

The parameters for D_T (and $D_T(|\mathbf{p}|, 0)$ itself), however, are strongly dependent on N_G if global updates are included. At $N_G = 100$ a plateau is not yet reached. In the deconfinement phase, results are also sensitive to the presence of global update steps, in particular at high values of β . Runs with only local updates produce propagators with significantly lower fit values of α_T and m_T , nearer to what is expected.

Realizing this, we decided to use in the final measurements at $T > 0$, in the region $\beta \geq 2.0$, only local updates and to use not less than $N_G = 100$ Gribov copies per configuration. Deep in confinement (below $\beta = 2.0$ for $T \neq 0$, and generally for $T = 0$) our standard was $N_G = 20$, with global updates included in the Monte Carlo. Thus, the results for D_T at larger β should be considered only as qualitative. The best fit parameters α and m of the propagators D , D_L and D_T are presented in Fig. 3 as functions of β . The anomalous dimensions $\alpha \neq 0$ in the confinement region for all propagators are functions not only of the monopole density (which is monotonously decreasing with growing β). The cluster structure of the monopole configurations seems to play a significant role. The fit parameters α_L and m_L of D_L vanish at β_c , giving a clear signal of the finite temperature phase transition [4] caused by the formation of dipoles. This is not the case for α_T and m_T of the D_T propagator. We associate the (so far) inconclusive behaviour of D_T , at the transition and beyond, with the insufficient gauge fixing. The mass m for $T = 0$ is in good agreement with the theoretical prediction [1] valid for a dilute monopole gas.

In conclusion, we have studied the gauge boson propagator in cQED₃, both at zero and non-zero temperatures. We have found that the propagators in all cases under investigation can be fitted by (3) which is a sum of the massive propagator with an anomalous dimension plus a contact




Figure 3. Fitted α and m vs. β for D , D_L , D_T .

term. Similar to D_L in the confinement phase [4], D for $T = 0$ has an anomalous dimension $\alpha \neq 0$. The behavior of D_T at the phase transition is obscured by the extreme sensitivity with respect to Gribov copy effects, *i.e.* remaining wrapping Dirac strings.

REFERENCES

1. A. M. Polyakov, Nucl. Phys. **B 120**, 429 (1977).
2. H. R. Fiebig and R. M. Woloshyn, Phys. Rev. **D 42**, 3520 (1990).
3. M. N. Chernodub, E.-M. Ilgenfritz, A. Schiller, Phys. Rev. **D 64**, 054507 (2001); *ibid.* 114502 (2001).
4. M. N. Chernodub, E.-M. Ilgenfritz, A. Schiller, Phys. Rev. Lett. **88**, 231601 (2002).
5. M. N. Chernodub, E.-M. Ilgenfritz, A. Schiller, hep-lat/0208013.
6. R. J. Wensley and J. D. Stack, Phys. Rev. Lett. **63**, 1764 (1989).
7. T. DeGrand and D. Toussaint, Phys. Rev. **D 22**, 2478 (1980).